

<<中国植物志>>

图书基本信息

书名 : <<中国植物志>>

13位ISBN编号 : 9781930723917

10位ISBN编号 : 1930723911

出版时间 : 2010-2

出版时间 : 科学出版社

作者 : Flora of China Editorial Committee 等著

页数 : 642

版权说明 : 本站所提供之下载的PDF图书仅提供预览和简介,请支持正版图书。

更多资源请访问 : <http://www.tushu007.com>

<<中国植物志>>

前言

Volume 10 of the Flora of China is the 18th of the 24 text volumes in the series. It includes one family , 29 tribes , 167 genera , and 1 , 673 species , among which one genus and 690 species (41%) are endemic to China , and three or four tribes , 32 or 33 genera , and 131-134 species (8%) are introduced to China.

Volume 10 is formatted in the same manner as the first 17 volumes of the series : vol. 17 (1994) , vol. 16 (1995) , vol. 15 (1996) , vol. 18 (1998) , vol. 4 (1999) , vol. 24 (2000) , vol. 8 (2001) , vol. 6 (2001) , vol. 9 (2003) , vol. 5 (2003) , vol. 14 (2005) , vol. 22 (2006) , vol. 13 (2007) , vol. 12 (2007) , vol. 11 (2008) , vol. 7 (2008) , and vol. 25 (2009) . The following 16 accompanying volumes of the Flora of China Illustrations have been published to date : vol. 17 (1998) , vol. 16 (1999) , vol. 15 (2000) , vol. 18 (2000) , vol. 4 (2001) , vol. 24 (2002) , vol. 8 (2003) , vol. 6 (2003) , vol. 9 (2004) , vol. 5 (2004) , vol. 14 (2006) , vol. 22 (2007) , vol. 13 (2008) , vol. 12 (2008) , vol. 11 (2009) , and vol. 7 (2009) . Bruce Bartholomew co-authored and edited the Hedysareae (Eversmannia) , Robinieae , and Sesbanieae and edited the Aeschynomeneae , Crotalarieae , Galegeae (Oxytropis) , Hedysareae (ex-cept , Alhagi , Calophaea , and Halimodendron) , and Millettiaeae. Anthony R. Brach co-authored and edited the Galegeae (Gueldenstaedtia and Tibetia) and edited the Acacieae , Caesalpinieae , Cassieae , Cercideae , Dalbergieae , Desmodieae , Detarieae , Euchrestaeae , Galegeae (except As-tragalus , Oxytropis , and Phyllobodium) , Hedysareae (Alhagi , Calophaea , and Halimodendron) , Ingeae , and Mimosaeae. Michael G. Gilbert co-authored and edited the Abreaeae , Amorpheae , and Psoraleeae , edited and partly co-authored the Phaseoleae , and edited the Indigofereae. Lisa J. Pepper and Nicholas J. Turland edited the Fabeae (Lathyrus) , Galegeae (Astragalus and Phyl-lolobium) , Loteae , Sophoreae , Trifolieae , and Thermopsideae.

<<中国植物志>>

内容概要

This volume entirely comprises the legume family (Fabaceae or Leguminosae), with 1,673 species in China, of which 690 are endemic, i.e., found nowhere else on Earth. The legumes of China are divided into 29 tribes, starting with the Cercideae, which include the disjunct redbuds (*Cercis*) and pantropical *Bauhinia*, and finishing with the Fabeae, which include vetches (*Vicia*), vetchlings (*Lathyrus*), lentil (*Lens culinaris*), and pea (*Pisum sativum*). The largest legume genus in China is *Astragalus*, with 401 species (it is also the largest genus globally, with ca. 3,000 species), followed by *Oxytropis* (133 species), *Indigofera* (79 species), in which *I. tinctoria* and *I. suffruticosa* are sources of the dye indigo, and *Caragana* (66 species).

<<中国植物志>>

书籍目录

Preface
Introduction
Acknowledgments
Fabaceae 豆科
dou ke
List of Nomenclatural Novelties Published in this Volume of the Flora of China
Index to Chinese Names
Index to Pinyin Names
Index to Scientific Names
Index to Families in the Flora of China and the Flora Reipublicae Popularis Sinicae
Published Volumes of the Flora of China and the Flora of China Illustrations

<<中国植物志>>

章节摘录

Lianas, woody, with tendrils; whole plant glabrous except for pedicel base and leaf buds. Twigs slightly zigzag, slender; tendrils solitary, slender. Leaves 2-foliate; stipules caducous; petiole 1-2 cm, slender, glabrous; leaf blade 12-24 x 9-16 mm, membranous, abaxially thinly pubescent on axis of primary veins, adaxially glabrous, veins 3 per lobe, veins dense, salient on both surfaces, base truncate; lobes obliquely obovate, apex obtuse or rounded. Inflorescence corymbose-racemose, terminal on lateral branches, many flowered; peduncle more than 1 cm; bracts and bracteoles linear, 1.5-4 mm. Pedicel 1-3 cm, slender. Flower buds ellipsoid, ca. 5 mm. Receptacle cylindric, 10-12 mm; calyx lobes 5, reflexed, broadly ovate or lanceolate. Petals white, broadly obovate, ca. 9 ~ 6 mm, abaxially hairy near base, shortly clawed. Fertile stamens 3; filaments ca. 1 cm, glabrous. Staminodes 3-5. Ovary shortly stalked, glabrous; style short; stigma disciform. Legume strap-shaped, compressed, ca. 10x 2.5 cm, tardily dehiscent, glabrous, suture slightly thickened. Seeds ellipsoid, compressed, ca. 6 x 4 mm; testa brown. Fl. Aug-Dec, fr. Sep-Feb.

<<中国植物志>>

媒体关注与评论

"...a landmark series of volumes" ——Economic Botany "...a must for any student of the temperate and subtropical flora of such a vast. species-rich country.." ——Kew Bulletin "...an indispensable tool for identifying Chineseplants..." ——Systematic Botany

<<中国植物志>>

版权说明

本站所提供下载的PDF图书仅提供预览和简介，请支持正版图书。

更多资源请访问:<http://www.tushu007.com>