

<<甚高能宇宙伽马辐射>>

图书基本信息

书名：<<甚高能宇宙伽马辐射>>

13位ISBN编号：9787301215531

10位ISBN编号：7301215533

出版时间：2012-12

出版时间：北京大学出版社

作者：阿哈隆尼安

页数：495

版权说明：本站所提供下载的PDF图书仅提供预览和简介，请支持正版图书。

更多资源请访问：<http://www.tushu007.com>

<<甚高能宇宙伽马辐射>>

内容概要

《中外物理学精品书系·引进系列(7)·甚高能宇宙伽马辐射：研究极端宇宙的关键窗口(影印版)》综述了宇宙伽马辐射这一现象的实验观测结果、起源的解释和蕴含的信息。其中主要探讨了宇宙射线的产生，相对论性射束的物理和天体物理，以及观测伽马射线宇宙学等研究方向。

具体包括银河系内和银河系外的宇宙射线的起源、极端天体物理条件(比如正在吸积的黑洞附近)下的加速和辐射过程、伽马射线暴等瞬态现象，以及与之相关的宇宙学课题等。

《中外物理学精品书系·引进系列(7)·甚高能宇宙伽马辐射：研究极端宇宙的关键窗口(影印版)》适合天文学、宇宙学、粒子物理等领域的研究者和研究生阅读。

<<甚高能宇宙伽马辐射>>

作者简介

作者：（德国）阿哈隆尼安（Aharonian F.A.）

书籍目录

Preface 1. Introduction 1.1 "The Last Electromagnetic Window" 1.2 Energy Domains of Gamma Ray Astronomy 1.3 Gamma Ray Astronomy: A Discipline in Its Own Right 2. Status of the Field 2.1 Low Energy Gamma Ray Sources 2.1.1 The COMPTEL source catalog 2.2 High Energy Gamma Ray Sources 2.2.1 GeV blazars 2.2.2 GeV pulsars 2.2.3 Unidentified EGRET sources 2.3 The Status of Ground-Based Gamma Ray Astronomy 2.3.1 Brief historical review 2.3.2 Reported TeV sources 2.3.2.1 The Crab Nebula 2.3.2.2 Other plerions 2.3.2.3 Gamma ray pulsars 2.3.2.4 Gamma rays from supernova remnants 2.3.2.5 Other galactic sources 2.3.2.6 TeV blazars 2.3.2.7 Other extragalactic objects 2.3.3 Next generation of IACT arrays Very High Energy Cosmic Gamma Radiation 2.3.3.1 Atmospheric Cherenkov radiation 2.3.3.2 Stereoscopic detection of Cherenkov images 2.3.3.3 IACT arrays 2.3.3.4 Sub-10 GeV ground based detectors? 2.3.3.5 Large field-of-view detectors 2.3.3.6 IACT arrays for probing PeV γ -rays 3. Gamma Ray Production and Absorption Mechanisms 3.1 Interactions with Matter 3.1.1 Electron bremsstrahlung and pair-production 3.1.2 Electron-positron annihilation 3.1.3 Gamma rays produced by relativistic protons 3.1.3.1 π^0 -decay gamma rays 3.1.3.2 Nuclear gamma ray line emission 3.2 Interactions with Photon Fields 3.2.1 Inverse Compton scattering 3.2.2 Photon-photon pair production 3.2.3 Interactions of hadrons with radiation fields 3.3 Interactions with Magnetic Fields 3.3.1 Synchrotron radiation and pair-production 3.3.2 Synchrotron radiation of protons 3.4 Relativistic Electron-Photon Cascades 4. Gamma Rays and Origin of Galactic Cosmic Rays 4.1 Origin of Galactic Cosmic Rays: General Remarks 4.1.1 What do we know about Cosmic Rays? 4.1.2 What we do not know about Cosmic Rays? 4.1.3 Common beliefs and "nasty" problems 4.1.4 Searching for sites of production of GCRs 4.2 Giant Molecular Clouds as Tracers of Cosmic Ray 4.2.1 Proton fluxes in the ISM near the accelerator 4.2.1.1 Impulsive source 4.2.1.2 Continuous source 4.2.1.3 The case of dense gas regions 4.2.2 Gamma rays from a cloud near the accelerator 4.2.3 Accelerator inside the cloud 4.2.4 On the level of the "sea" of galactic cosmic rays 4.3 Probing the Sources of VHE CR Electrons 4.3.1 Distributions of VHE electrons 4.3.2 Extended regions of IC gamma radiation 4.4 Diffuse Radiation from the Galactic Disk 4.4.1 CR spectra in the inner Galaxy 4.4.2 Diffuse radiation associated with cosmic ray electrons 4.4.2.1 IC gamma rays 4.4.2.2 Electron bremsstrahlung 4.4.2.3 Annihilation of CR positrons in flight 4.4.3 Gamma rays of nucleonic origin 4.4.4 Overall gamma ray fluxes 4.4.5 Probing the diffuse γ -ray background on small scales 4.4.6 Concluding remarks 5. Gamma Ray Visibility of Supernova Remnants 5.1 Gamma Rays as a Diagnostic Tool 5.2 Inverse Compton Versus π^0 -Decay Gamma Rays 5.3 Synchrotron X-ray Emission of SNRs 5.4 TeV Gamma Radiation of SN 1006 and Similar SNR 5.4.1 Inverse Compton models of TeV emission 5.4.2 Hadronic origin of TeV emission? 5.4.3 Distinct features of electronic and hadronic models 5.4.4 Concluding remarks 5.5 Molecular Clouds Overtaken by SNRs 5.5.1 Bremsstrahlung X-rays from Cygni 5.5.2 The case of RX J1713.7-3946 5.6 A Special Case: Gamma Rays from Cassiopeia A 5.7 "PeV SNRs" 6. Pulsars, Pulsar Winds, Plerions 6.1 Magnetospheric Gamma Rays 6.1.1 Polar cap versus outer gap models 6.1.2 Magnetospheric TeV gamma rays? 6.2 Gamma Rays from Unshocked Pulsar Winds 6.2.1 Characteristics of the KED wind 6.2.2 The ejection rate and the particle spectrum 6.2.3 IC Radiation of the pulsar wind in Crab 6.2.4 Gamma rays from winds of PSR B1706-44 and Vela? 6.2.5 IC γ -rays from the binary pulsar PSR B1259-63 6.3 Gamma Rays from Pulsar Driven Nebulae 6.3.1 Broad-band nonthermal radiation of the Crab Nebula 6.3.1.1 Synchrotron and IC radiation 6.3.1.2 Second High Energy Synchrotron Component 6.3.1.3 Bremsstrahlung and π^0 -decay gamma rays? 6.3.1.4 The objectives of future gamma ray studies 6.4 High Energy Gamma Rays from Other Plerions 6.4.1 Time-evolution of electrons 6.4.2 Target photon fields 6.4.3 Effects of B-field, electron energy, and pulsar age 6.4.4 Synchrotron and IC nebulae around PSR B1706-44 7. Gamma Rays Expected from Microquasars 7.1 Do We Expect Gamma Rays from X-Ray Binaries? 7.2 Nonthermal Phenomena in Microquasars 7.3 Modelling of Radio Flares of GRS 1915+105 7.4 Expected Gamma Ray Fluxes 7.5 Searching for Gamma Ray Signals from Microquasars 7.6 The Case of Microblazars 7.7 Ultraluminous Sources as Microblazars? 7.8 Persistent Gamma Ray Emission from Extended Lobes 8. Large Scale Jets of Radio Galaxies and Quasars 8.1 Synchrotron and IC Models of Large Scale AGN Jets 9. Nonthermal Phenomena in Clusters of Galaxies 10. TeV Blazars and Cosmic Background

<<甚高能宇宙伽马辐射>>

Radiation 11. High Energy Gamma Rays - Carriers of Unique Cosmological Information Appendix A Spherically symmetric diffusion from a single source Appendix B Evolution of relativistic electrons in an expandid magnetised medium B.1 Kinetic equation B.2 Time-independent energy losses B.3 Expanding cloud Bibliography Index

<<甚高能宇宙伽马辐射>>

章节摘录

版权页：插图： On theoretical grounds, the diffusive shock acceleration model faces several challenges or "nasty problems" (Drury et al., 2001) like the "injection problem" and the "maximum energy problem", recently critically reviewed by Kirk and Dendy (2001), Drury (2001) and Malkov and Drury (2001). Diffusive shock acceleration requires particles with energy at least several times larger than the thermal energy of the plasma, and it is not yet clear how to get particles from the thermal pool accelerated to supra-thermal energies. Recent theoretical progress in this direction (e.g. Malkov and Sisk, 1995; Dieckmann et al., 2000) provides optimism that eventually the injection problem will be resolved, most likely through extensive numerical simulations (Kirk and Dendy, 2001). The problem of the maximum achievable energy problem is an old one and has a vital implication for the SNR paradigm of GCRs. In diffusive shock acceleration theory, the maximum energy of particles is achieved during the so-called free-expansion phase which, however, does not last long enough to allow acceleration of particles up to the highly desired point, the knee around 10¹⁵ eV. Therefore, violation of the so-called "upper limit" of Lagage and Cesarsky (1983), which, for the standard SNR parameters, the shock speed, duration of the free-expansion phase, and the ambient magnetic field, cannot significantly exceed 10¹⁴ eV, remains as one of the highest priorities of current theoretical studies. A promising way has recently been suggested by Lucek and Bell (2000). They showed that cosmic ray streaming drives large-amplitude Alfvénic waves which may amplify the magnetic field non-linearly to many times the pre-shock value. Thus, the cosmic rays themselves provide the field necessary for their effective acceleration! The increased magnetic field reduces the acceleration time, and correspondingly increases the maximum particle energies to 10¹⁵ eV and even beyond.

<<甚高能宇宙伽马辐射>>

编辑推荐

《甚高能宇宙伽马辐射:研究极端宇宙的关键窗口(影印版)》适合天文学、宇宙学、粒子物理等领域的研究者和研究生阅读。

<<甚高能宇宙伽马辐射>>

版权说明

本站所提供下载的PDF图书仅提供预览和简介，请支持正版图书。

更多资源请访问:<http://www.tushu007.com>